



## CELLULITE: EVALUATION AND TREATMENT

J.P. ORTONNE

Dermatology Department, Hôpital Archet, Nice France

Cellulite, described first in 1904, is a female feature which induces obsessional suffering in 30% of the women. Two hypothesis exist to describe this phenomenon depending where the anomalies are considered first: hypodermis and derm structure<sup>1</sup> or blood and lymphatic microvessels<sup>2</sup>. Whatever the initial mechanism, this disorder is associated with adipocyte hypertrophy (herniation of the dermo hypodermal junction) and circulatory deteriorations. Different methods can be used to study both components. In addition to classical methods such as photograder, standardized photographs, circumference and cutaneous fold measurements, more specialized techniques are now available to determine the efficacy of anti-cellulite treatment. Photogrammetry has shown that dietary complement containing amongst others ingredients bioflavonoids and ginko biloba induce a significant slimming effect<sup>3</sup>. Sonography, much more specific for the evaluation of skin structure itself, has been mainly used to test efficacy of topical treatments containing herbals extracts<sup>4</sup>, retinol<sup>5</sup> or retinol associated with caffeine and ruscogenin<sup>6</sup>. The different results are more or less convincing. Among the different treatments described in literature, Endermologie® caught our attention. Thanks to a range of advanced investigation methods (histochemistry, image analysis, laser doppler, sonography 20 Mz, prints analysis, macrophotographies), this technique has really proven its efficacy in cellulite linked to its dermatrophic, circulatory and anti edematous properties<sup>7,8,9,10</sup>.

In conclusion, either cellulite or anti-cellulite treatments must be evaluated preferably with combination of appropriate techniques in order to assess reliable data.

### REFERENCES

1. Nurnberger F, Mullr G. So-called cellulite : an invented disease. *J Dermatol Surg Oncol* 1978 ; 4 : 221-9.
2. Merlen JF, Curri SB, Sarteel AM. La cellulite, affection microvasculoconjunctive. *Phlébologie* 1979; 32 (3) : 279-82.
3. Soler-Rouanet 2001 Etude par photogrammétrie de l'efficacité amincissante du complément Cellulase Urto chez des volontaires sains féminins
4. Bascaglia DA, Conte ET, McCain W., Frideman S. The treatment of cellulite with methylxanthine and herbal extract based cream : an ultrasonographic analysis. *Cosmet Dermatol* 1996 ; 9 : 30-40.
5. Piérard Franchimont et al. 2000. A randomized, placebo-controlled trial of topical retinol in the treatment of cellulite
6. Bertin et al. 2001. A double-blind evaluation of the activity of an anti-cellulite product containing retinol, caffeine and ruscogenine by a combination of several non-invasive methods.
7. Adcock D., Paulsen S., Jabour K., Davis S., Nanney LB., Bruce Shack R. Analysis of the Effects of Deep Mechanical Massage in the Porcine Model. *Plast. Reconstr. Surg.* 2001 Jul., 108 (1); 233-40.
8. Watson J. , Fodor PB., Cutcliffe B., Sayah D., Shaw W. Physiological effects of Endermologie® : A Preliminary Report. *Aesthetic Surg J* 1999, 19 (1) ; 27-33.
9. Innocenzi D., Balzani A., Montesi G., La Torre G., Tenna S., Scuderi N., Calvieri S. Evidenza delle modificazioni cutanee indotte dalla Tecnica LPG® mediante analisi d'immagine . *DermoCosmetologia*-anno II- numero 1- gennaio-marzo 2003, p.9-15.
10. Innocenzi D., Balzani A., Panetta C., Montesi G., Tenna S., Scuderi N., Calvieri S. Alterazioni morfologiche della cute indotte con la Tecnica LPG®. *DERMOtime* Settembre-Ottobre 2002, anno XIV, n° 7/8 ; p. 25-27.